

FIRST RESULTS FROM PHENIX AT RHIC

A. BAZILEVSKY for the PHENIX Collaboration
RIKEN BNL Research Center, Brookhaven National Laboratory
Upton, NY 11973-5000, USA

The first results from Au-Au collisions at $\sqrt{s_{NN}}=130$ GeV obtained with the PHENIX detector in the Year 2000 run at RHIC are presented. The mid-rapidity charged particle multiplicity and transverse energy per participating nucleon rise steadily with the number of participants, such that $\langle E_T \rangle / \langle N_{ch} \rangle$ remains relatively constant as a function of centrality. Identified charged hadron spectra as well as $\bar{p}/p$ and K^+/K^- ratios are discussed. Charged particle and neutral pion transverse momentum distributions in peripheral nuclear collisions are consistent with point-like scaling. The spectra at high p_t from central collisions are significantly suppressed when compared to a simple superposition of binary nucleon-nucleon collisions.

1 Introduction

The Relativistic Heavy-Ion Collider (RHIC) at Brookhaven National Laboratory started regular operation in June 2000, opening new frontiers in the study of hadronic matter under unprecedented conditions of temperature and energy density.

PHENIX¹ is one of the four experiments at RHIC. It is designed to simultaneously measure a wide range of probes – photons, leptons and hadrons – sensitive to all timescales, from initial hard scattering to final state interactions.

All results presented in this paper are obtained using the two PHENIX central arm spectrometers covering approximately ± 0.35 in pseudorapidity and π in azimuth. A highly segmented Electromagnetic calorimeter (EMCal) allowed measurements of photons and neutral pions and also provided global event characterization through transverse energy. Charged particles were reconstructed using the Drift Chambers (DC), Pad Chambers (PC) and Time-of-Flight (ToF) walls. The momentum resolution achieved so far with DC is $\sigma_p/p^2 = 3.5\%/(GeV/c)$ for particles with momentum above 1 GeV/c. Particle identification was performed by ToF measurements. A large Ring Imaging Cherenkov counter (RICH) was used for electron identification.

Two other detectors, Beam-Beam Counters (BBC) and Zero Degree Calorimeters (ZDC), provided a trigger and the off-line event selection². The BBC comprise two arrays of 64 quartz radiator Cherenkov detectors, located at ± 1.44 m from the center of the interaction region (IR) covering 2π in ϕ and $3.0 < |\eta| < 3.9$. The BBC also provide the start timing and vertex position³. The ZDC are small transverse-area hadron calorimeters that measure neutron energy within $|\eta| > 6$. The centrality bins were determined based upon the BBC vs. ZDC signal distribution. Simulations of the response of the BBC and the ZDC were used to account for the effect of physics and detector fluctuations in the definition of these event classes and to relate them via a Glauber model to the number of participating nucleons N_p and the number of binary collisions N_c ².




Figure 1: Multiplicity (left) and Transverse energy (right) densities per participant pair at mid-rapidity as function of the number of participants.

2 Charged particle multiplicity and transverse energy

The number of primary charged particles per event was determined on a statistical basis by correlating two layers of Pad Chambers placed at radial distances of 2.49 m and 4.98 m, respectively, from the IR². A section of PbSc Calorimeter provided the transverse energy measurements⁴. Fig.1 shows the results for the multiplicity and transverse energy densities per pair of participating nucleons as a function of N_p . Both quantities exhibit very consistent behavior, such that transverse energy per charged particle, $\langle E_T \rangle / \langle N_{ch} \rangle$, remains relatively constant at a value ~ 0.8 GeV (Fig.2 left). The same value for the most central collisions can be derived from the measurements done by WA98⁵ and NA49⁶ at SPS at $\sqrt{s_{NN}}=17.2$ GeV and from the measurement of E814/E877⁷ at AGS at $\sqrt{s_{NN}}=4.8$ GeV (Fig.2 right). Coming back to Fig.1, it should be noted that the slope parameters α in $dN_{ch}/d\eta$ and $dE_T/d\eta$ parameterizations $\propto N_p^\alpha$ obtained at RHIC^{2,4,8} are larger than at SPS: our data give $\alpha = 1.16 \pm 0.04$ ($\alpha = 1.13 \pm 0.05$) for multiplicity density (transverse energy density) compared to $\alpha = 1.07 \pm 0.04$ ($\alpha = 1.08 \pm 0.06$) found by the WA98 experiment⁵. That allows us to state that at RHIC energy the “wounded nucleon model” ($\alpha = 1$) breaks down and a term proportional to the number of collisions makes a significant contribution to particle and energy production and it is larger than at SPS where α is found to be close to unity within the errors of measurements. Fig.1 (left) also shows $dN_{ch}/d\eta|_{\eta=0}$ for $N_p = 2$ from UA5¹⁵ data for $p\bar{p}$.

Our straightforward derivation of Bjorken energy density ϵ_{Bj} from our measured $dE_T/d\eta|_{\eta=0}$ (and corrected to $dE_T/dy|_{y=0}$) for the most central 2% of the inelastic cross section gives $\epsilon_{Bj} = 4.6$ GeV/fm³ (ref.⁴), an increase of more than 60% over the NA49 value⁹.




Figure 2: Transverse energy per charged particle at mid-rapidity vs N_{part} (left) and energy of participating nucleons (right) for the highest centrality bin.




Figure 3: Minimum bias transverse momentum distribution for positive (left) and negative (right) identified hadrons. The error bars include statistical errors and systematic errors in the acceptance and decay corrections. Additional 20% systematic errors on the absolute normalization are not included.

3 Hadron yields

The ToF time resolution of 115 ps allowed pion and kaon separation up to 1.5–2 GeV/c, and proton separation up to 3.5 GeV/c. Fig.3 shows the minimum bias p_t spectra for $\pi^{+/-}$, $K^{+/-}$, p and $\bar{p}$. The inverse slope parameters increase with particle mass¹⁰, such that protons and antiprotons become the significant contributors to hadron p_t spectra above 2 GeV/c.

The $\bar{p}/p$ and K^+/K^- ratios for minimum bias collisions was found to be $0.64 \pm 0.01(stat) \pm 0.07(syst)$ and $1.08 \pm 0.03(stat) \pm 0.22(syst)$ correspondingly. Within uncertainties, neither transverse momentum dependence nor centrality dependence were seen in the $\bar{p}/p$ and K^+/K^- ratios¹¹. At RHIC energies both $\bar{p}/p$ and K^+/K^- ratios are clearly approaching unity, indicating that pair creation of $\bar{p}/p$ and K^+/K^- in the midrapidity region is becoming a dominant process and implying that the net baryon density is much less than that measured in lower energy experiments at AGS^{12,13} and SPS¹⁴.

4 High transverse momentum particles

In heavy ion collisions at RHIC energies jet production is expected to dominate particle production at high p_t . In the high density medium created in central heavy ion collisions it is predicted that scattered partons may lose considerable energy via gluon bremsstrahlung or “jet quenching”^{16,17}. It may result in a reduction of the high p_t hadron yield.

Fig.4 shows the transverse momentum distribution of π^0 's in peripheral (upper 60–80% of σ_{int}) and the 10% most central Au-Au collisions. Both spectra are compared to different theoretical calculations¹⁶. The peripheral data are consistent with pQCD calculations for pp , with simple point-like scaling to Au-Au collisions by the mean number of binary collisions. By contrast, the central data points are well below curves calculated without accounting for parton energy loss¹⁸. The significant suppression of high p_t particles in central collisions was also observed in charged hadron spectra¹⁹.




Figure 4: Comparison of π^0 spectra to theoretical calculations under three scenarios and for two centralities. The points are PHENIX data. The curves are calculations of X-N. Wang: solid lines are a pQCD calculation for pp scaled by the number of binary collisions; the dotted lines add shadowing and p_t broadening; the dashed lines add a $dE/dx = 0.25$ GeV/fm parton energy loss.

5 Conclusion

In just a few months after the first Au-Au collisions were observed at RHIC in June 2000, the PHENIX collaboration produced the first intriguing physics results. Because of the limited size of the paper we presented here only some of the PHENIX results regarding charged multiplicity and transverse energy distributions, hadron yields and spectra of high transverse momentum particles. One of the unique PHENIX capabilities is electron identification in a wide momentum range. The first inclusive electron spectra and electron pair mass distribution were obtained from year 2000 data²⁰. In the coming run in 2001, PHENIX will accumulate data of about three orders of magnitude larger, so that we expect to observe several thousand $J/\psi, \phi \rightarrow e^+e^-$ events.

Acknowledgments

We thank the staff of the RHIC project, Collider-Accelerator, and Physics Departments at BNL and the staff of PHENIX participating institutions for their vital contributions. We acknowledge support from the Department of Energy and NSF (U.S.A.), Monbu-sho and STA (Japan), RAS, RMAE, and RMS (Russia), BMBF and DAAD (Germany), FRN, NFR, and the Wallenberg Foundation (Sweden), MIST and NSERC (Canada), CNPq and FAPESP (Brazil), IN2P3/CNRS (France), DAE (India), KRF and KOSEF (Korea), and the US-Israel Binational Science Foundation.

References

1. PHENIX Collaboration, D. P. Morrison, *et al.*, Nucl. Phys. **A 638**, 565 (1998).
2. PHENIX Collaboration, K. Adcox, *et al.*, Phys. Rev. Lett. **86**, 3500 (2001).
3. K. Ikematsu *et al.*, Nucl. Instrum. Meth. **A411**, 238 (1998).
4. PHENIX Collaboration, K. Adcox, *et al.*, nucl-ex/0104015; submitted to Phys. Rev. Lett.
5. WA98 Collaboration, M. M. Aggarwal, *et al.*, Eur. Phys. J. **C18**, 651 (2001).

6. NA49 Collaboration, T. Alber, *et al.*, *Phys. Rev. Lett.* **75**, 3814 (1995).
7. E814/E877 Collaboration, J. Barrette, *et al.*, *Phys. Rev. Lett.* **70**, 2996 (1993).
8. A. Milov, for the PHENIX Collaboration, to be published in the Proceedings of Quark Matter 2001 Conference.
9. NA49 Collaboration, T. Alber, *et al.*, *Phys. Rev. Lett.* **75**, 3814 (1995). (The quoted $\epsilon_{Bj} = 3.2 \text{ GeV/fm}^3$ was divided by 1.10 to remove the enhancement factor for “head-on” collisions).
10. J. Velkovska, for the PHENIX Collaboration, to be published in the Proceedings of Quark Matter 2001 Conference, nucl-ex/0105012.
11. H. Ohnishi, for the PHENIX Collaboration, to be published in the Proceedings of Quark Matter 2001 Conference.
12. L. Ahle, *et al.*, *Phys. Rev. C* **58**, 3523 (1998).
13. L. Ahle, *et al.*, *Phys. Rev. Lett.* **81**, 2650 (1998).
14. M. Kaneta, *et al.*, *J.Phys.G: Nucl. Part. Phys.* **23** (1997) 1865.
15. G.J. Alner *et al.*, *Z. Phys.* **C33**, 1 (1986).
16. X.N. Wang, *Phys. Rev. C* **61**, 064910 (2000).
17. P. Levai, G. Papp, G. Fai, M. Gyulassy, nucl-ex/0104015.
18. G. David, for the PHENIX Collaboration, to be published in the Proceedings of Quark Matter 2001 Conference, nucl-ex/0105014.
19. F. Messer, for the PHENIX Collaboration, to be published in the Proceedings of Quark Matter 2001 Conference.
20. Y. Akiba, for the PHENIX Collaboration, to be published in the Proceedings of Quark Matter 2001 Conference.